






Annual Examinations for Secondary Schools 2022


YEAR 9	CHEMISTRY	MARKING SCHEME
---------------	------------------	-----------------------

Item			Answer and guidelines	Marks
Section A				
1	a		Evaporation.	1
	b	i	T	1
		ii	F	1
		iii	T	1
		iv	F	1
	c	i	F	1
		ii	T	1
		iii	F	1
		iv	F	1
	d	i	chemical	1
		ii	physical	1
		iii	physical	1
			Total	12
2	a		b	1
	b		Only one type of atom makes up iodine molecules.	1
	c	i	Any one of: <ul style="list-style-type: none"> • Shiny when freshly cut • Good conductors of heat and electricity • Relatively high melting point • Relatively high density • Malleable (can be hammered) • Ductile (can be drawn into wires) • Usually solid at room temperature • Most metals are sonorous or make a bell-like sound when struck <i>Accept other suitable answers.</i>	1
		ii	Made of more than one type of atom chemically combined.	1
		iii	$\text{Fe} + \text{S} \rightarrow \text{FeS}$ <i>1 mark for formulae, 1 mark for balancing.</i>	2
			Total	6
3	a		carbon dioxide; water	1,1
	b	i	Presence of oxygen or air; presence of water	1,1
		ii	A	1
		iii	Nail in test-tube A is in water and exposed to oxygen in air.	1
			Total	6

Item			Answer and guidelines	Marks
4	a		 <p>1 mark for drawing atoms joined correctly. 1 mark for showing correctly shared electrons. 1 mark for correct number of electrons. Accept atoms touching each other.</p>	3
	b		 <p>1 mark for each ion drawn correctly including brackets, 1 mark for correct charges.</p>	3
			Total	6
5	a	i	C	1
		ii	A	1
	b	i	increases	1
		ii	decreases	1
	c		<p>1 kg = 1000 g</p> <p>In 1000 g water, 5000 g sugar dissolve</p> <p>In 100 g water, ?</p> <p>$5000 \times 100 / 1000 = 500 \text{ g/100 g H}_2\text{O}$</p>	1
			Total	6
6	a		Sample B	1
			Sample A	1
	b		<p>A: Temporary and permanent hardness.</p> <p>B: Temporary hardness.</p> <p><i>Do not award marks if for A, only one type of hardness is given.</i></p>	1
	c		Permanent hardness	1
	d		On adding soap solution, soft water lathers while hard water does not.	1
			Total	6
7	a	i	Orange juice	1
		ii	Does not ionise completely in solution so it gives fewer H^+ ions.	1
		iii	Tonic water.	1
	b	i	<p>$\underline{1}\text{H}_2\text{SO}_4(\text{aq}) + \underline{2}\text{NaOH}(\text{aq}) \rightarrow \underline{1}\text{Na}_2\text{SO}_4(\text{aq}) + \underline{2}\text{H}_2\text{O}(\text{l})$</p> <p>1 mark for formulae, 1 mark for balancing. Accept as correct if no number is given where balancing is 1.</p>	2
		ii	<p>$\underline{2}\text{HNO}_3(\text{aq}) + \underline{\text{K}_2}\text{CO}_3(\text{aq}) \rightarrow \underline{2}\text{KNO}_3(\text{aq}) + \underline{\text{H}_2}\text{O}(\text{l}) + \underline{1}\text{CO}_2(\text{g})$</p> <p>1 mark for formulae, 1 mark for balancing. Accept as correct if no number is given where balancing is 1.</p>	2
	c		Sodium hydrogen sulfate - NaHSO_4	1
	d	i	Lead(II) nitrate; sodium chloride	1,1

Item			Answer and guidelines	Marks
			<i>Accept any other suitable soluble salts.</i>	
		ii	$\text{Pb}(\text{NO}_3)_2 + 2\text{NaCl} \rightarrow \text{PbCl}_2 + 2\text{NaNO}_3$ <i>1 mark for formulae, 1 mark for balancing.</i>	2
			Total	12
8	a	i	RFM of NH_4NO_3 : $(14 \times 2) + (1 \times 4) + (3 \times 16) = 80$	1
		ii	% nitrogen by mass: $(28/80)100 = 35\%$	1
	b	i	Divide by RAM: Carbon: $83.7/12 = 6.975$ Hydrogen: $(100-83.7)/1 = 16.300$ Divide by smallest to work out ratio: Carbon: $6.975/6.975 = 1.000$ Hydrogen: $16.300/6.975 = 2.337$ Simplest ratio (multiply by 3): Carbon: $1.000 \times 3 = 3.000$ Hydrogen: $2.337 \times 3 = 7.011$ Empirical formula: C_3H_7 <i>Apply follow through.</i>	1 1 1
		ii	RMM of $\text{C}_3\text{H}_7 = (12 \times 3) + (1 \times 7) = 43$ $86/43 = 2$ Therefore, molecular formula is $(\text{C}_3\text{H}_7) \times 2 = \text{C}_6\text{H}_{14}$ <i>Apply follow through.</i>	1
			Total	6
Section B				
9	a	i	Method: <ul style="list-style-type: none"> Add water and stir well. Filter mixture. Insoluble silver chloride is obtained as residue. Silver nitrate solution is obtained as filtrate. Wash residue well with distilled water to obtain silver chloride in pure form. Leave it to dry. Evaporate the water slowly from the silver nitrate solution. Use a water bath to heat the solution (avoid direct heating since silver nitrate decomposes on heating). 	1 1 1 1 1
		ii	Filtration:  <i>1 mark for filter funnel.</i> <i>1 mark for beaker or conical flask.</i> <i>1 mark for correctly labelling silver chloride residue.</i> <i>1 mark for clearly labelling silver nitrate filtrate solution.</i> <i>Deduct ½ mark for each missing label.</i>	4

Item			Answer and guidelines	Marks
			Evaporation:  <p> 1 mark for evaporating dish. 1 mark for drawing water bath. 1 mark for drawing tripod & Bunsen burner. 1 mark for clearly labelling silver nitrate solution. Deduct ½ mark for each missing label. Award only 1 mark if the direct heated method is chosen and drawn correctly. </p>	4
	b	i	Fractional distillation.	1
		ii	Cools down any vapour passing through it which turns to liquid.	1
		iii	Improves the separation of liquids with different boiling points being distilled.	1
		iv	Water	1
		v	Liquids have different boiling points. The liquid with the lower boiling point boils off first.	1 1
		vi	Fractional distillation of crude oil <i>Accept fractional distillation of liquid air.</i>	1
			Total	20
10	a	i	$2\text{H}_2\text{O}_2(\text{aq}) \rightarrow 2\text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$ <i>1 mark for formulae, 1 mark for balancing, 1 mark for state symbols.</i>	3
		ii	Manganese(IV) oxide / MnO_2	1
		iii	 <p> 1 mark each for reaction flask, dropping funnel. Accept any suitable type of flask. 2 marks for gas collection apparatus: gas jar, trough & shelf. Accept any suitable gas collection apparatus. 1 mark for correct labelling of reagents used. Deduct ½ mark for each missing label. </p>	5
	b	i	$5.1 - 2.4 = 2.7 \text{ g H}_2\text{O}$	1

		ii	$\text{RMM H}_2\text{O}_2 = (1 \times 2) + (16 \times 2) = 34$ $34.0 \text{ g} : 1 \text{ mole}$ $5.1 \text{ g} : ?$	1
			$5.1/34 = \mathbf{0.15 \text{ moles H}_2\text{O}_2}$ <i>Allow follow through.</i>	1
	c	i	Fe_2O_3 : basic Al_2O_3 : amphoteric CO: neutral SO_2 : acidic	1 1 1 1
		ii	$\text{RFM of Fe}_2\text{O}_3 = (56 \times 2) + (16 \times 3) = \mathbf{160}$ $\text{RMM of CO}_2 = 12 + (16 \times 2) = \mathbf{44}$ Mole ratio $\text{Fe}_2\text{O}_3 : \text{CO}_2 = \mathbf{1 : 3}$ Therefore $160 \text{ g Fe}_2\text{O}_3 : (44 \times 3) = 132 \text{ g CO}_2$ $160,000 \text{ g} : (132 \times 160,000) / 160 = \mathbf{132000 \text{ g CO}_2}$ Moles $\text{CO}_2 = 132000 / 44 = \mathbf{3000 \text{ moles}}$ <i>Accept if moles of CO₂ are calculated before mass.</i> <i>Allow follow through.</i>	1 1 1 1
			Total	20
11	a	i	$2\text{HCl(aq)} + \text{Zn(s)} \rightarrow \text{ZnCl}_2\text{(aq)} + \text{H}_2\text{(g)}$ <i>1 mark for formulae, 1 mark for balancing, 1 mark for state symbols.</i> <i>Accept other suitable metals.</i>	3
		ii	 <i>1 mark each for reaction flask, dropping funnel.</i> <i>Accept any suitable type of flask/dropping or thistle funnel.</i> <i>2 marks for gas collection apparatus: gas jar, trough & shelf.</i> <i>Accept any suitable gas collection apparatus.</i> <i>1 mark for correct labelling of reactants.</i> <i>Deduct ½ mark for each missing label.</i>	5
		iii	Insert a lighted wooden splint into a test tube full of hydrogen. Hydrogen burns with a 'pop'.	1 1

	b		$9.2\text{ g} + 0.4\text{ g} = \mathbf{9.6\text{ g}}$	1
	c	i	$\text{CuO} + \text{H}_2 \rightarrow \text{Cu} + \text{H}_2\text{O}$ <i>1 mark for formulae, 1 mark for balancing.</i>	2
		ii	RFM of CuO = $63.5 + 16 = 79.5$ Moles of CuO in 159 g = $159/79.5 = \mathbf{2\text{ moles}}$ 1 mole of CuO contains $6.02 \times 10^{23} \times 2 = 1.204 \times 10^{24}$ ions (since each CuO has 2 ions (Cu^{2+} and O^{2-})) So, 2 moles CuO have = $1.204 \times 10^{24} \times 2 = \mathbf{2.408 \times 10^{24}\text{ ions}}$	1 1 1
	d		RFM $\text{Fe}_2\text{O}_3 = (56 \times 2) + (16 \times 3) = \mathbf{160}$ Mole ratio $\text{Fe}_2\text{O}_3 : \text{Fe} = \mathbf{1 : 2}$ Therefore 160 g $\text{Fe}_2\text{O}_3 : (56 \times 2) = 112\text{ g Fe}$ 1600 g : $(112 \times 1600) / 160 = \mathbf{1120\text{ g Fe}}$ Moles Fe = $1120 / 56 = \mathbf{20\text{ moles}}$ <i>Accept if moles of Fe are calculated before mass.</i>	1 1 1 1
			Total	20